



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Combined agonist-antagonist genome-wide functional screening identifies broadly active antiviral microRNAs

Citation for published version:

Santhakumar, D, Forster, T, Laqtom, NN, Fragkoudis, R, Dickinson, P, Abreu-Goodger, C, Manakov, SA, Choudhury, NR, Griffiths, SJ, Vermeulen, A, Enright, AJ, Dutia, B, Kohl, A, Ghazal, P & Buck, AH 2010, 'Combined agonist-antagonist genome-wide functional screening identifies broadly active antiviral microRNAs', *Proceedings of the National Academy of Sciences (PNAS)*, vol. 107, no. 31, pp. 13830-13835. <https://doi.org/10.1073/pnas.1008861107>

Digital Object Identifier (DOI):

[10.1073/pnas.1008861107](https://doi.org/10.1073/pnas.1008861107)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Proceedings of the National Academy of Sciences (PNAS)

Publisher Rights Statement:

PNAS open access option

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Combined agonist–antagonist genome-wide functional screening identifies broadly active antiviral microRNAs

Diwakar Santhakumar^{a,b}, Thorsten Forster^{b,c}, Nouf N. Laqtom^{a,b,d}, Rennos Fragkoudis^e, Paul Dickinson^{b,c},
Cei Abreu-Goodger^f, Sergei A. Manakov^f, Nila Roy Choudhury^e, Samantha J. Griffiths^b, Annaleen Vermeulen^g,
Anton J. Enright^f, Bernadette Dutia^e, Alain Kohl^e, Peter Ghazal^{b,c}, and Amy H. Buck^{a,b,1}

^aCentre for Immunity, Infection and Evolution, University of Edinburgh, Edinburgh EH9 3JT, United Kingdom; ^bDivision of Pathway Medicine and Centre for Infectious Diseases, University of Edinburgh, Edinburgh EH16 4SB, United Kingdom; ^cCentre for Systems Biology at Edinburgh, University of Edinburgh, Edinburgh EH9 3JD, United Kingdom; ^dDepartment of Biology, King Abdulaziz University, Jeddah 21589, Saudi Arabia; ^eThe Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh EH9 1QH, United Kingdom; ^fEuropean Bioinformatics Institute, Cambridge CB10 1SD, United Kingdom; and ^gThermo Fisher Scientific, Dharmacon Products, Lafayette, CO 80026

Communicated by Norman R. Pace, University of Colorado, Boulder, CO, June 28, 2010 (received for review April 26, 2010)

Although the functional parameters of microRNAs (miRNAs) have been explored in some depth, the roles of these molecules in viral infections remain elusive. Here we report a general method for global analysis of miRNA function that compares the significance of both overexpressing and inhibiting each mouse miRNA on the growth properties of different viruses. Our comparative analysis of representatives of all three herpesvirus subfamilies identified host miRNAs with broad anti- and proviral properties which extend to a single-stranded RNA virus. Specifically, we demonstrate the broad antiviral capacity of miR-199a-3p and illustrate that this individual host-encoded miRNA regulates multiple pathways required and/or activated by viruses, including PI3K/AKT and ERK/MAPK signaling, oxidative stress signaling, and prostaglandin synthesis. Global miRNA expression analysis further demonstrated that the miR-199a/miR-214 cluster is down-regulated in both murine and human cytomegalovirus infection and manifests similar antiviral properties in mouse and human cells. Overall, we report a general strategy for examining the contributions of individual host miRNAs in viral infection and provide evidence that these molecules confer broad inhibitory potential against multiple viruses.

RNAi | herpesvirus | RNA virus | RNA processing | phosphatidylinositol-3-kinase-Akt signalling

Since the discovery of the first microRNA (miRNA) in *Caenorhabditis elegans*, research in diverse organisms has illuminated the role of this class of small RNA in a wide range of cellular processes (reviewed in ref. 1). MicroRNAs modulate the expression of specific genes by guiding the RNA-induced silencing complex (RISC) to complementary sites within messenger RNAs (mRNAs) (2). This generally serves to down-regulate target genes at specific times, in concert with other regulatory mechanisms in the cell (reviewed in ref. 3). Functional analysis of individual miRNAs suggests diversity in the timing and mechanisms by which they regulate cellular events. For example, some miRNAs promote cellular proliferation, whereas others promote apoptosis (depending on which genes are targeted). Consequently, the expression level of a given miRNA within a cell is expected to be under tight regulatory control, and mechanisms for achieving this control continue to emerge (reviewed in ref. 4).

Viruses require host-cell processes for their survival and have evolved mechanisms for modifying cellular conditions toward an environment conducive to replication while evading recognition and destruction by the host. Herpesviruses are one of the oldest and most successful viral families in this regard. They have coevolved with their hosts for hundreds of millions of years and infect nearly all vertebrate species studied and at least one invertebrate (5). Human cytomegalovirus (HCMV), a member of the β -herpesvirus subfamily that infects a large portion of the world's population (50–90%), is a leading cause of congenital infection (~1% of live births) and a major cause of morbidity in immunocompromised patients. Although mouse and human cytomegaloviruses have diverged over ~80 million years, the

pathophysiology of murine CMV (MCMV) in mice is similar to that of HCMV in humans, and the lytic infections result in activation and manipulation of common host-cell signaling cascades (6).

Because miRNAs regulate many aspects of cellular physiology, their expression levels could impact the infection process. It might be expected, therefore, that host miRNA expression is subject to regulation upon infection, by either viral or host factors. Indeed, we and others have previously identified host miRNAs that are down-regulated upon infection by cytomegaloviruses (in some cases within 4 h postinfection) and have demonstrated that these miRNAs exert antiviral properties when overexpressed (7, 8). However, to date, there has been no overlap between results reported with MCMV and HCMV, nor has there been any context with which to interpret the significance of the effect of overexpressing a given miRNA in relation to any other miRNA in the cell. Similarly, various groups have identified mammalian miRNAs that are regulated or implicated in diverse viral infections, but the majority of these studies are generally founded on expression profiling or miRNA target predictions, (reviewed in ref. 9; ref. 10). There is relatively little investigation to date on the functional impact of miRNAs in different infections.

We postulate that specific subsets of host miRNAs are important in controlling the infection process and might be subject to regulation by host and/or viral factors. Although the kinetic parameters of miRNA action are not well-known (e.g., how quickly and reversibly they can modulate a gene or pathway), we expect that viruses with slower replication kinetics (>24 h) might be particularly sensitive to (and exploitive of) changes in host miRNA expression levels. Here we report a combined agonist–antagonist miRNA screening approach that is designed to obtain functional information about mouse miRNAs that impact the lytic phase of herpesviral infections. We further test the breadth of observed antiviral miRNA properties in human cells and against an evolutionarily unrelated RNA virus. Our functional and expression analyses demonstrate that host miRNAs are a tunable and important component of herpesviral infection and provide evidence that these molecules have broad antiviral properties in mouse and human cells against both DNA and RNA viruses.

Author contributions: D.S., N.N.L., A.K., and A.H.B. designed research; D.S., N.N.L., R.F., and A.H.B. performed research; N.R.C., S.J.G., A.V., B.D., A.K., and P.G. contributed new reagents/analytic tools; D.S., T.F., N.N.L., P.D., C.A.-G., S.A.M., A.J.E., and A.H.B. analyzed data; and A.H.B. wrote the paper.

Conflict of interest statement: A.V. is, as of the publication date, employed, with fixed salaries, by Thermo Fisher Scientific, which offers for sale libraries of miRNA mimics and inhibitors.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: a.buck@ed.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1008861107/-DCSupplemental.




Fig. 4. Changes in miRNA expression upon MCMV or HCMV infection. (A) Heat map displaying differential expression of mouse and human miRNAs in MCMV (NIH 3T3 cells, MOI = 3; 24 hpi) or HCMV (MRC-5 cells, MOI = 3; 48 hpi). Color scale is based on log-2 changes in expression; probes identified as significant in each dataset are bracketed (left, MCMV; right, HCMV), with overlap depicted. FC, fold change. (B) Northern blot analysis of total RNA isolated from mouse fibroblast NIH 3T3 cells infected with MCMV at MOI = 3 (24 hpi) or human fibroblast MRC-5 cells infected with HCMV at MOI = 3 (48 hpi). Dashed lines indicate a gap between lanes that was removed in making the image.

Kaposi's sarcoma-associated herpesvirus infection that facilitates viral replication and regulates antiviral innate immunity (28). This further supports the concept that some mammalian hosts and


viruses might have evolved mechanisms to regulate individual miRNAs (or specific subsets) upon infection, rather than targeting the global miRNA machinery. Further work in other infection systems and cell types is required for a more global perspective of the antiviral miRNAs identified here, as well as for miR-132, which was excluded from our analysis due to nonspecific effects in the mimic library (*SI Methods*).

We previously reported rapid down-regulation of miR-27a and miR-27b in MCMV infection and speculated that this regulation might reflect antiviral functions of these miRNAs (8). Neither miR-27a nor miR-27b is among the most significant antiviral miRNAs identified here. However, our analysis is limited to those miRNAs that display the most dramatic (and opposite) effects on viral growth when overexpressed and inhibited in fibroblasts. Although this allows for high confidence in the hits identified, it may miss more subtle, but real, effects of other miRNAs and will be cell-type-specific. An underlying assumption is also that overexpression of a miRNA increases the extent to which it represses its targets; this depends on the concentrations of miRNAs and targets, which will be variable in different cells. Further understanding of miRNA biology is required to shed light on the generality of this assumption. Although miR-27a and miR-27b are not scored as significant in the mimic libraries (according to rank product analysis), they are among the most significant antiviral miRNAs when examining the combined inhibitor datasets for all three viruses, along with miR-24 and miR-23b (*Dataset S1*), with which they are coregulated.

We focus here on miRNAs with common properties in multiple herpesviral infections; however, there are notable differences in the screening results with different viruses (for example, miR-378 is proviral in MCMV but not the other infections; Fig. 2B). Viruses can evolve miRNA binding sites in their genomes (29), and it is possible that differences observed here are related to sequence-specific miRNA interactions with viral elements. Given the expected frequency of miRNA target sites in a ~200,000-base-pair herpesviral genome (on average, a 7-mer seed site will occur every ~16,384 nt), investigations in this area will require extensive biochemical validation.

To date, the therapeutic capacity of miRNA manipulation in viral infection has focused primarily on blocking interactions between a host miRNA and a viral sequence (for example, miRNA-122/HCV) (29, 30). Here we find that miR-199a-3p leads to decreased viral growth in all three herpesviral subfamilies as well as an unrelated positive-strand RNA virus. Although we cannot eliminate the possibility that the antiviral properties are at least partly

Fig. 5. Cellular networks modulated by miR-199a-3p. (A) Plot of the significance of enrichment of all 7-mer words complementary to known mouse miRNA seed regions in the mRNA array dataset, comparing mimic against inhibitor results. The x axis represents all of the genes with an annotated 3'UTR sequence, sorted starting from the most down-regulated by the miR-199a-3p mimic (equivalent to the most up-regulated by the inhibitor). The y axis represents the $-\log_{10}$ transformed *P* values, with positive values denoting enrichment and negative values denoting depletion. Each gray line represents a single 7-mer, tested for enrichment and depletion at every 250 genes. Two seed-matching 7-mers for miR-199a-3p, 7-mer(1A) = CTACTGA and 7-mer(2) = ACTACTG, are highlighted in gold and red, respectively. Horizontal dotted lines represent Bonferroni-corrected *P* value significance thresholds of 0.01. (B) Pathways regulated by miR-199a-3p that are known to play a role in CMV infection. Selected genes down-regulated by miR-199a-3p are shown in blue circles (and are shaded if they also represent a predicted target); human alias gene names are used.



based on interactions with viral elements [as suggested in HCV (31)], we identify several host pathways regulated by this miRNA that are activated and required by multiple viruses. Specifically, ERK/MAPK signaling, prostaglandin synthesis, oxidative stress signaling, and PI3K/AKT signaling are all down-regulated by miR-199a-3p (Fig. 5 and Table S1). The regulation of these pathways by miR-199a-3p is consistent with previous (although diverse) reports on miR-199a-3p, a stress-inducible miRNA up-regulated during cardiac hypertrophy (32) that is also reported to target MET (33), a transmembrane tyrosine kinase receptor associated with invasive growth of tumors, and COX2 (34), an enzyme involved in prostaglandin synthesis. Our findings suggest links between this miRNA and viral infection. Greater understanding of the role of miR-199a-3p in normal and disease biology will shed light on the best strategies for exploiting its antiviral properties. It is notable that the other miRNA with which it is coregulated, miR-214, is among the four miRNAs identified here as broadly antiviral, (Figs. 2 and 3) and may function in concert with miR-199a-3p.

Given the large number of potential targets of any given miRNA [on average, in the hundreds (35)], and in particular those described in this study, it may not be that one specific target (or

even a handful of targets) sufficiently explains a miRNA-related phenotype. We anticipate that understanding miRNA function may require a “whole is greater than the sum of the parts” approach (involving multiple targets), which might also hold true in developing more effective antiviral therapeutics.

Methods

For a full description of the methods used here, see *SI Methods*. The miRNA mimic and inhibitor libraries are based on miRBase 8.2. The engineered reporter viruses have been described elsewhere (14, 36–39). MicroRNA microarrays are based on Exiqon miCURY probe set version 8.1. Pathway analysis was carried out using Ingenuity software and targets were predicted based on targetscan.org (TargetScanHuman version 5.1) (20).

ACKNOWLEDGMENTS. We thank Giles Dudley and Jon Karpilow for helpful discussions related to this work and Marie Craigon, David Page, and Alan Ross at the Division of Pathway Medicine for technical assistance. This work is supported by Biotechnology and Biological Sciences Research Council (BBSRC) Grant BB/G01552X/1 and Wellcome Trust CIE Grant 082611/Z/07/Z (A.H.B), Wellcome Trust Programme Grant WT 066784/Z/02/Z and the Centre for Systems Biology at Edinburgh funded by BBSRC and the Engineering and Physical Sciences Research Council (EPSRC), reference BB/D019621/1 (P.G.), and BBSRC Roslin Institute Strategic Programme Grant (A.K.).

- Chang TC, Mendell JT (2007) microRNAs in vertebrate physiology and human disease. *Annu Rev Genomics Hum Genet* 8:215–239.
- Lee RC, Feinbaum RL, Ambros V (1993) The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 75:843–854.
- Bartel DP (2009) MicroRNAs: Target recognition and regulatory functions. *Cell* 136: 215–233.
- Winter J, Jung S, Keller S, Gregory RI, Diederichs S (2009) Many roads to maturity: MicroRNA biogenesis pathways and their regulation. *Nat Cell Biol* 11:228–234.
- Davison AJ (2002) Evolution of the herpesviruses. *Vet Microbiol* 86:69–88.
- Krmpotic A, Bubic I, Polic B, Lucin P, Jonjic S (2003) Pathogenesis of murine cytomegalovirus infection. *Microbes Infect* 5:1263–1277.
- Wang FZ, et al. (2008) Human cytomegalovirus infection alters the expression of cellular microRNA species that affect its replication. *J Virol* 82:9065–9074.
- Buck AH, et al. (2010) Post-transcriptional regulation of miR-27 in murine cytomegalovirus infection. *RNA* 16:307–315.
- Umbach JL, Cullen BR (2009) The role of RNAi and microRNAs in animal virus replication and antiviral immunity. *Genes Dev* 23:1151–1164.
- Parameswaran P, et al. (2010) Six RNA viruses and forty-one hosts: Viral small RNAs and modulation of small RNA repertoires in vertebrate and invertebrate systems. *PLoS Pathog* 6:e1000764.
- Vermeulen A, et al. (2007) Double-stranded regions are essential design components of potent inhibitors of RISC function. *RNA* 13:723–730.
- Birmingham A, et al. (2009) Statistical methods for analysis of high-throughput RNA interference screens. *Nat Methods* 6:569–575.
- Breitling R, Armengaud P, Amtmann A, Herzyk P (2004) Rank products: A simple, yet powerful, new method to detect differentially regulated genes in replicated microarray experiments. *FEBS Lett* 573:83–92.
- Kiiver K, et al. (2008) Properties of non-structural protein 1 of Semliki Forest virus and its interference with virus replication. *J Gen Virol* 89:1457–1466.
- Mankouri J, et al. (2010) Optineurin negatively regulates the induction of IFN β in response to RNA virus infection. *PLoS Pathog* 6:e1000778.
- Baek D, et al. (2008) The impact of microRNAs on protein output. *Nature* 455:64–71.
- Selbach M, et al. (2008) Widespread changes in protein synthesis induced by microRNAs. *Nature* 455:58–63.
- van Dongen S, Abreu-Goodger C, Enright AJ (2008) Detecting microRNA binding and siRNA off-target effects from expression data. *Nat Methods* 5:1023–1025.
- Fortunato EA, McElroy AK, Sanchez I, Spector DH (2000) Exploitation of cellular signaling and regulatory pathways by human cytomegalovirus. *Trends Microbiol* 8: 111–119.
- Lewis BP, Burge CB, Bartel DP (2005) Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120: 15–20.
- Feire AL, Koss H, Compton T (2004) Cellular integrins function as entry receptors for human cytomegalovirus via a highly conserved disintegrin-like domain. *Proc Natl Acad Sci USA* 101:15470–15475.
- Johnson RA, Wang X, Ma XL, Huang SM, Huang ES (2001) Human cytomegalovirus up-regulates the phosphatidylinositol 3-kinase (PI3-K) pathway: Inhibition of PI3-K activity inhibits viral replication and virus-induced signaling. *J Virol* 75:6022–6032.
- Johnson RA, Ma XL, Yurochko AD, Huang ES (2001) The role of MKK1/2 kinase activity in human cytomegalovirus infection. *J Gen Virol* 82:493–497.
- Schröer J, Shenk T (2008) Inhibition of cyclooxygenase activity blocks cell-to-cell spread of human cytomegalovirus. *Proc Natl Acad Sci USA* 105:19468–19473.
- Zhu H, Cong JP, Yu D, Bresnahan WA, Shenk TE (2002) Inhibition of cyclooxygenase 2 blocks human cytomegalovirus replication. *Proc Natl Acad Sci USA* 99:3932–3937.
- Otsuka M, et al. (2007) Hypersusceptibility to vesicular stomatitis virus infection in Dicer1-deficient mice is due to impaired miR24 and miR93 expression. *Immunity* 27: 123–134.
- Pfeffer S, et al. (2004) Identification of virus-encoded microRNAs. *Science* 304: 734–736.
- Lagos D, et al. (2010) miR-132 regulates antiviral innate immunity through suppression of the p300 transcriptional co-activator. *Nat Cell Biol* 12:513–519.
- Jopling CL, Yi M, Lancaster AM, Lemon SM, Sarnow P (2005) Modulation of hepatitis C virus RNA abundance by a liver-specific microRNA. *Science* 309:1577–1581.
- Lanford RE, et al. (2010) Therapeutic silencing of microRNA-122 in primates with chronic hepatitis C virus infection. *Science* 327:198–201.
- Murakami Y, Aly HH, Tajima A, Inoue I, Shimotohno K (2009) Regulation of the hepatitis C virus genome replication by miR-199a. *J Hepatol* 50:453–460.
- van Rooij E, et al. (2006) A signature pattern of stress-responsive microRNAs that can evoke cardiac hypertrophy and heart failure. *Proc Natl Acad Sci USA* 103: 18255–18260.
- Kim S, et al. (2008) MicroRNA miR-199a* regulates the MET proto-oncogene and the downstream extracellular signal-regulated kinase 2 (ERK2). *J Biol Chem* 283: 18158–18166.
- Chakrabarty A, et al. (2007) MicroRNA regulation of cyclooxygenase-2 during embryo implantation. *Proc Natl Acad Sci USA* 104:15144–15149.
- Friedman RC, Farh KK, Burge CB, Bartel DP (2009) Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res* 19:92–105.
- Dutia BM, et al. (2004) Identification of a region of the virus genome involved in murine herpesvirus 68-induced splenic pathology. *J Gen Virol* 85:1393–1400.
- Sun A, et al. (2004) Immediate-early expression of the herpes simplex virus type 1 ICP27 transcript is not critical for efficient replication in vitro or in vivo. *J Virol* 78: 10470–10478.
- Angulo A, et al. (2000) Identification of a boundary domain adjacent to the potent human cytomegalovirus enhancer that represses transcription of the divergent UL127 promoter. *J Virol* 74:2826–2839.
- Angulo A, Ghazal P, Messerle M (2000) The major immediate-early gene ie3 of mouse cytomegalovirus is essential for viral growth. *J Virol* 74:11129–11136.